

AM I AT RISK FOR TYPE 2 DIABETES?

Taking Steps To Lower the Risk of Getting Diabetes



National Institutes of Health National Institute of Diabetes and Digestive and Kidney Diseases

Table of Contents

What is diabetes?
How can type 2 diabetes be prevented?
What are the signs and symptoms of type 2 diabetes? 2
Should I be tested for diabetes?4
What does it mean to be at risk for type 2 diabetes? 4
What are my risk factors for developing type 2 diabetes?5
What can I do about my risk factors? 6
Doing my part: Getting started
Hope through research

What is diabetes?

Diabetes is a disease in which blood glucose levels are above normal. People with diabetes have problems converting food to energy. After a meal, food is broken down into a sugar called glucose, which is carried by the blood to cells throughout the body. Cells use a chemical known as insulin, made in the pancreas, to help them process blood glucose into energy.

People develop diabetes for two reasons: the pancreas does not make enough insulin for the body's needs, or the cells in the muscles, liver, and fat do not use insulin properly, or both. As a result, the amount of glucose in the blood increases while the cells are starved of energy. Over the years, high blood glucose damages nerves and blood vessels, leading to complications such as blindness, heart and kidney disease, nerve problems, gum infections, and amputation.

Although people with diabetes can prevent or delay complications by keeping blood glucose levels close to normal, preventing or delaying the development of diabetes in the first place is even

better. The results of the Diabetes Prevention Program (DPP) show how to do so.

How can type 2 diabetes be prevented?

A recently concluded Federally funded study of 3,234 people at high risk for diabetes, the Diabetes Prevention Program, showed that diet and exercise can sharply lower your risk of getting type 2 diabetes.

The DPP was a major clinical trial that studied ways to prevent or delay diabetes

in people at high risk for type 2 diabetes. Participants were overweight and had higher than normal levels of blood glucose, called impaired glucose tolerance. Both conditions are strong risk factors for type 2 diabetes. Because of the high risk among some minority groups, about half of the DPP participants were African American, American Indian, Asian American, Pacific Islander, or Hispanic.

The DPP compared two approaches to preventing diabetes: an intensive healthy eating and exercise program and the diabetes drug metformin. People who engaged in moderate physical activity for about 30 minutes a day, followed a low-fat, low-calorie diet, and lost 5 to 7 percent of their body weight reduced their risk of getting type 2 diabetes by 58 percent, according to preliminary analysis of the results. Those receiving metformin reduced their risk by 31 percent.

What are the signs and symptoms of type 2 diabetes?

Many people have no signs or symptoms. Symptoms can also be so mild that you might not even notice them. Five million people in the United States have type 2 diabetes and do not know it. Here is what to look for:

- increased thirst
- increased hunger
- fatigue
- increased urination, especially at night
- weight loss
- blurred vision
- sores that do not heal

Types of diabetes

The three main kinds of diabetes are type 1, type 2, and gestational diabetes.



Type 1 diabetes

Type 1 diabetes, formerly called juvenile diabetes or insulindependent diabetes, is usually first diagnosed in children, teenagers, or young adults. In this form of diabetes, the beta cells of the pancreas no longer make insulin because the body's immune system has attacked and destroyed them. Treatment for type 1 diabetes includes taking insulin shots or using an insulin pump, making wise food choices, exercising regularly, taking aspirin daily, and controlling blood pressure and cholesterol.



Type 2 diabetes

Type 2 diabetes, formerly called adult-onset diabetes or noninsulin-dependent diabetes, is the most common form of diabetes. People can develop type 2 diabetes at any age—even during childhood. In type 2 diabetes, the pancreas does not make enough insulin and the fat, muscle, and liver cells do not use it properly. Being overweight and inactive can increase the chances of developing type 2 diabetes. Treatment includes taking diabetes medicines, making wise food choices, exercising regularly, taking aspirin daily, and controlling blood pressure and cholesterol.



Gestational diabetes

Some women develop gestational diabetes during the late stages of pregnancy. Although this form of diabetes usually goes away after the baby is born, a woman who has had it is more likely to develop type 2 diabetes later in life. Gestational diabetes is caused by the hormones of pregnancy or a shortage of insulin.

Sometimes people have symptoms but do not suspect diabetes. They delay scheduling a checkup because they do not feel sick. Many people do not find out they have the disease until they have diabetes complications, such as blurry vision or heart trouble. It is important to find out early if you have diabetes because treatment can prevent damage to the body from diabetes.

Should I be tested for diabetes?

If you are 45 years old or older, you need to get tested for diabetes, even if you do not have any of the risk factors listed on page 5. If you are younger than 45 and have one or more risk factors, you should also get tested. Ask your doctor for a fasting blood glucose test. The results of this blood test will tell your doctor how much glucose is in your blood. Your doctor may ask you to have the test twice.

If your blood glucose level is normal and you have no risk factors, you may need to remind your doctor to check your blood glucose again in 3 years. If you have at least one of the risk factors on page 5, have your blood checked once a year.

What does it mean to be at risk for type 2 diabetes?

You are more likely to get type 2 diabetes if you are age 45 or older, are overweight, are inactive, or have high blood pressure or high cholesterol. You are also at high risk if diabetes runs in your family or if you are African American, American Indian, Asian American, Hispanic, or Pacific Islander. However, you can take steps to reduce your risk of developing diabetes.

What are my risk factors for developing type 2 diabetes?

To find out your risk for type 2 diabetes, check each item that applies to you.

I have a parent, brother, or sister with diabetes.
My family background is African American, American Indian, Asian American, Hispanic, or Pacific Islander.
I have had gestational diabetes or I gave birth to at least one baby weighing 9 pounds or more.
I have been told that my blood glucose is higher than normal but not high enough to be diagnosed with diabetes. In other words, I have impaired glucose tolerance or an impaired fasting glucose level.
I am 45 years old or older.
I am overweight. My weight is at least 20 percent higher than it should be for my height. (See the chart on pages 10 and 11.)
I am fairly inactive. I exercise fewer than three times a week.
My blood pressure is 140/90 or higher, or I have been told that I have high blood pressure.
My cholesterol levels are not normal. My LDL cholesterol is higher than 100; my HDL cholesterol ("good" cholesterol) is 35 or lower; or my triglyceride level is 250 or higher.

If you checked one or more boxes, you are at risk for type 2 diabetes. If you checked more than three boxes, your chances of getting diabetes are high. It is important to lower your risk by working on the factors under your control.

What can I do about my risk factors?

You can do a lot to lower your chances of getting diabetes. Exercising regularly, reducing fat and calorie intake, and losing weight can all help you reduce your risk of developing type 2 diabetes. Lowering blood pressure and cholesterol levels also help you stay healthy.

If you checked

☐ I am overweight.

Then take these steps:

- Reach and maintain a reasonable body weight (see page 8).
- Make wise food choices most of the time (see page 9).
- Be physically active every day (see page 9).

If you checked

■ I am fairly inactive.

Then take this step:

• Be physically active every day (see page 9).

-If you checked

☐ My blood pressure is 140/90 or higher.

Then take these steps:

- Reach and maintain a reasonable body weight (see page 8).
- Make wise food choices most of the time (see page 9).
- Reduce your intake of salt and alcohol.
- Talk to your doctor about whether you need medicine to control your blood pressure (see page 12).

If you checked

☐ My cholesterol levels are not normal.

Then take these steps:

- Make wise food choices most of the time (see page 9).
- Be physically active every day (see page 9).
- Talk to your doctor about whether you need medicine to control your cholesterol levels (see page 12).

Doing my part: Getting started

Making big changes in your life is hard, especially if you are faced with more than one change. You can make it easier by taking these steps:

- Make a plan to change behavior.
- Decide exactly what you will do and when you will do it.
- Plan what you need to get ready.
- Think about what might prevent you from reaching your goals.
- Find family and friends who will support and encourage vou.
- Decide how you will reward yourself when you do what you have planned.

Your doctor, a dietitian, or a counselor can help you make a plan. Here are some of the areas you may wish to change to reduce your risk of diabetes.

Reach and maintain a reasonable body weight

Your weight affects your health in many ways. Being overweight can keep your body from making and using insulin properly. It can also cause high blood pressure. The DPP showed that losing even a few pounds can help reduce your risk of developing type 2 diabetes because it helps your body use insulin more effectively. In the DPP, people who lost between 5 and 7 percent of their body weight significantly reduced their risk of type 2 diabetes. For example, if you weigh 150 pounds, losing about 10 pounds would make a difference.

Body mass index (BMI) is a measure of body weight relative to height. You can use BMI to see whether you are underweight, normal weight, overweight, or obese. Use the table on pages 10 and 11 to find your BMI.

- Find your height in the left-hand column.
- Move across in the same row to the number closest to your weight.
- The number at the top of that column is your BMI. Check the word above your BMI to see whether you are underweight, normal weight, overweight, or obese.

If you are overweight or obese, choose sensible ways to get in shape:

- Avoid crash diets. Instead, eat less of the foods you usually have. Limit the amount of fat you eat.
- Increase your physical activity. Aim for at least 30 minutes of exercise most days of the week. (See page 9 for easy suggestions.)
- Set a reasonable weight-loss goal, such as losing 1 pound a
 week. Aim for a long-term goal of losing at least 7 percent
 of your total body weight.

Make wise food choices most of the time

What you eat has a big impact on your health. By making wise food choices, you can help control your body weight, blood pressure, and cholesterol.

- Take a hard look at the serving sizes of the foods you eat.
 Reduce serving sizes of main courses (such as meat),
 desserts, and foods high in fat. Increase the amount of fruits and vegetables.
- Limit your fat intake to about 25 percent of your total calories. For example, if your food choices add up to about 2,000 calories a day, try to eat no more than 56 grams of fat. Your doctor or a dietitian can help you figure out how much fat to have. You can check food labels for fat content too.
- You may also wish to reduce the number of calories you have each day. Your doctor or dietitian can help you with a meal plan that emphasizes weight loss.
- Keep a food and exercise log. Write down what you eat, how much you exercise— anything that helps keep you on track.
- When you meet your goal, reward yourself with a nonfood item or activity, like watching a movie.

Be physically active every day

Regular exercise tackles several risk factors at once. It helps you lose weight, keeps your cholesterol and blood pressure under control, and helps your body use insulin effectively. People in the DPP who were physically active for 30 minutes a day reduced their risk of type 2 diabetes. Many chose brisk walking for exercise.

If you are not very active, you should start slowly, talking with your doctor first about what kinds of exercise would be safe for you.

¥////	E E	40		TO THE OWNER OF THE OWNER OWN	5				E VE		列聚		375	Bod	N N	ass	<u>Ind</u>	ex	Body Mass Index Table	0	1	VA		132	11, E.S.			1/2/		Sin B		EMER		
			No	Normal				Over	Overweight	ŧ			Obese	ese									ш	xtre	ne O	Extreme Obesity	λ							
BMI	19	70	7	52	73	24 2	25 2	26 2	27 28	8 29	9 30	3	32	33	8	35	98	37	88	စ္တ	40	14	42	43 4	4 4	45 46	6 47	48	49	20	51	25	53	54
Height (inches)	<u> </u>														Bod	y Wei	Body Weight (pounds)	unoc	(sp															
28	91	96	100	100 105 110 115	110 1		119 1	24 1;	124 129 134 138 143	4 13	88 14	4	8 153	3 158	3 162	167	172		177 181	186	191	196	201 2	205 2	210 21	215 220	20 224	4 229	234	239	244	248	253 2	258
29	94	66	104	109	114 1	119 1	124 13	128 13	133 13	138 143	148	8 153	3 158	8 163	3 168	173	178	183	188	193	198	203	208 2	212 2	217 22	222 227	27 232	2 237	242	247	252	257	262	267
09	97	102	107	112	118 1	123 1	128 1:	133 13	138 14	143 148	R 153	3 158	8 163	3 168	3 174	179	184	189	194	199	204	209	215 2	220 2	225 23	230 235	35 240	0 245	250	255	261	266	271	276
61	100	106	11	116	122 1	127 1	132 1	137 14	143 148	153	3 158	8 164	4 169	9 174	180	185	190	195	201	206	211	217	222	227 2	232 23	238 243	13 248	8 254	259	264	269	275	280	285
62	104	109	115	120	126 1	131 1	136 1	142 14	147 153	3 158	164	4 169	9 175	5 180	186	191	196	202	207	213	218	224	229 2	235 2	240 24	246 251	51 256	5 262	267	273	278	284	289 2	295
63	107	113	118	124	130 1	135 1	141	146 1	152 158	58 163	33 169	9 175	5 180	0 186	191	197	203	208	214	220	225	231	237 2	242 2	248 25	254 259	59 265	5 270	278	282	287	293	299	304
64	110	116	122	128	134 1	140	145 1	151 18	157 16	163 169	99 174	9	0 186	6 192	197	204	. 209	215	221	227	232	238	244 2	250 2	256 26	262 267	37 273	3 279	285	291	296	302	308	314
65	114	120	126	132	138 1	4	150 1	156 16	162 16	168 174	74 180	0 186	6 192	2 198	3 204	210	216	222	228	234	240	246	252 2	258 2	264 27	270 276	76 282	2 288	294	300	306	312	318	324
99	118	124	130	136	142 1	148	155 1	161 16	167 17	173 179	79 186	6 192	2 198	8 204	1 210	216	223	229	235	241	247	253	260 2	266 2	272 27	278 284	34 291	1 297	303	309	315	322	328	334
29	121	127	134	140	146 1	153 1	159 10	166 17	172 17	178 185	35 191	1 198	8 204	4 211	217	, 223	230	236	242	249	255	261	268 2	274 2	280 28	287 293	33 299	908 6	312	319	325	331	338	344
89	125	131	138	44	151 1	158 1	164 1	171 17	177 184	34 190	197	7 203	3 210	0 216	3 223	230	236	243	249	256	262	769	276 2	282 2	289 29	295 302	308	3 315	322	328	335	341	348	354
69	128	135	142	149	155 1	162 1	169 1	176 18	182 189	39 196	96 203	3 209	9 216	6 223	3 230	236	243	250	257	263	270	277	284 2	291 2	297 30	304 311	11 318	324	331	338	345	351	358	365
20	132	139	146	153	160 1	167 1	174 18	181	188 195	35 202	209	9 216	6 222	2 229	236	243	250	257	264	271	278	285	292 2	299 3	306 31	313 320	20 327	7 334	341	348	355	362	369	376
71	136	143	150	157	165 1	172 1	179 18	186 19	193 200	00 208	215	22	2 229	9 236	3 243	250	257	265	272	279	786	293	301 3	308 3	315 32	322 329	29 338	343	351	358	365	372	379	386
72	140	147	154	. 791	169 1	177 1	184 1	191 18	199 206	06 213	13 221	1 228	8 235	5 242	250	258	265	272	279	287	294	302	309 3	316 3	324 33	331 338	38 346	353	361	368	375	383	330	397
73	144	151	159	166	174 1	182 1	189 1	197 20	204 212	12 219	19 227	7 235	5 242	2 250) 257	, 265	272	280	288	295	302	310	318 3	325 3	333 34	340 348	18 355	5 363	371	378	386	393	401 4	408
74	148	155	163	171	179 1	186 1	194 20	202 27	210 218	18 225	25 233	3 24	1 249	9 256	3 264	272	280	287	295	303	311	319	326 3	334 3	342 35	350 358	365	5 373	381	389	396	404	412 4	420
75	152	160	168	176	184 1	192 2	200 2	208 27	216 224	24 232	32 240	0 248	8 256	6 264	1 272	279	287	295	303	311	319	327	335 3	343 3	351 35	359 367	37 375	5 383	391	333	407	415	423 4	431
9/	156	164	172	180	189 1	197 2	205 2	213 22	221 230	30 238	38 246	22	4 263	3 271	279	287	295	304	312	320	328	336	344 3	353 3	361 36	369 377	77 385	5 394	402	410	418	426	435 4	443
																															ı	١	ı	

Source: Adapted from Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults: The Evidence Report.

Make a plan to increase your activity level toward the goal of being active for at least 30 minutes a day most days of the week.

Choose activities you enjoy. Here are some ways to work extra activity into your daily routine:

- Take the stairs rather than an elevator or escalator.
- · Park at the far end of the lot and walk.
- Get off the bus a few stops early and walk the rest of the way.
- Walk or bicycle instead of drive whenever you can.

Take your prescribed medications

Some people need medication to help control their blood pressure or cholesterol levels. If you do, take your medicines as directed. Ask your doctor whether there are any medicines you can take to prevent type 2 diabetes.

Hope through research

We now know that regular exercise, weight loss, and a low-fat diet can prevent type 2 diabetes. However, scientists are still investigating whether type 1 diabetes can be prevented.

In the Diabetes Prevention Trial—Type 1, a study sponsored by the National Institute of Diabetes and Digestive and Kidney Diseases, close relatives of people with type 1 diabetes are tested for their risk of getting the disease. Those at high risk will receive treatment that might prevent type 1 diabetes. More information about the study is available by calling 1-800-425-8361 or at www.niddk.nih.gov/patient/dpt_1/dpt_1.htm.

Other studies on preventing type 1 diabetes are under way. For upto-date information on clinical trials, see *http://clinicaltrials.gov.*

National Diabetes Information Clearinghouse

1 Information Way Bethesda, MD 20892-3560

Phone: 1-800-860-8747 or (301) 654-3327

Fax: (301) 907-8906

Email: ndic@info.niddk.nih.gov

The National Diabetes Information Clearinghouse (NDIC) is a service of the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). The NIDDK is part of the National Institutes of Health under the U.S. Department of Health and Human Services. Established in 1978, the clearinghouse provides information about diabetes to people with diabetes and to their families, health care professionals, and the public. NDIC answers inquiries, develops and distributes publications, and works closely with professional and patient organizations and Government agencies to coordinate resources about diabetes.

Publications produced by the clearinghouse are carefully reviewed by both NIDDK scientists and outside experts. This booklet was reviewed by David G. Marrero, Ph.D., Indiana University School of Medicine, Diabetes Research and Training Center; and Michael L. Parchman, M.D., M.P.H., Associate Professor, Department of Family and Community Medicine, University of Texas Health Sciences Center.

This publication is not copyrighted. The clearinghouse encourages users of this booklet to duplicate and distribute as many copies as desired.

This booklet is also available at www.niddk.nih.gov under "Health Information."



National Institute of Diabetes and Digestive and Kidney Diseases NIH Publication No. 02–4805 November 2001